

YOUR HEALTHCARE PROVIDER HAS PRESCRIBED RELISTOR® FOR YOU...

Many patients who take chronic opioid pain medication may become constipated.

<https://www.relistor.com>

For more information and resources, scan the QR code below or go to www.Relistor.com



~80% of patients taking daily oral opioid therapy and OTC laxatives reported still being constipated.

RELISTOR works differently from over-the-counter (OTC) laxatives. Some common OTC laxatives include: Miralax®, Colace®, and Senokot®.* RELISTOR is an FDA-approved prescription treatment specifically designed to target the constipating effect opioids have in the gut, without impacting opioid pain relief. Tell your doctor if you experience any changes in your pain relief.

*All trademarks are the properties of their respective owners.



HOW TO TAKE RELISTOR TABLETS

RELISTOR tablets should be taken **DAILY**, as directed by your healthcare provider.



Take three 150 mg Relistor tablets, a total of 450 mg with water in the morning at least 30 minutes prior to your first meal of the day.

RELISTOR should be taken **DAILY** as long as you are taking opioid pain medication. Your doctor may change your dose if you have kidney or liver problems.



You don't have to accept substitutions from the pharmacy - your healthcare provider specifically prescribed RELISTOR for you. If needed, follow up with your doctor's office for additional assistance.

INDICATIONS

RELISTOR (methylnaltrexone bromide) is a prescription medicine used to treat constipation in adults that is caused by prescription pain medicines called opioids.

- RELISTOR tablets are used to treat constipation caused by opioids in adults with long-lasting (chronic) pain that is not caused by active cancer.

It is not known if RELISTOR is safe and effective in children.

IMPORTANT SAFETY INFORMATION

- Do not take RELISTOR if you have a bowel blockage (called an intestinal obstruction) or have a history of bowel blockage.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Medication Guide.

RELISTOR®
methylnaltrexone bromide
Tablets

IMPORTANT SAFETY INFORMATION (Continued)

- RELISTOR can cause serious side effects such as a tear in your stomach or intestinal wall (perforation). Stomach pain that is severe can be a sign of a serious medical condition. If you get stomach pain that is severe, does not go away, or gets worse, stop taking RELISTOR and get emergency medical help right away.
- Stop using RELISTOR and call your healthcare provider if you get diarrhea that is severe or that does not go away during treatment with RELISTOR.
- You may have symptoms of opioid withdrawal during treatment with RELISTOR including sweating, chills, diarrhea, stomach pain, anxiety, and yawning. Tell your healthcare provider if you have any of these symptoms.
- Tell your healthcare provider if you have kidney or liver problems.
- Tell your healthcare provider if you have any stomach or bowel (intestines) problems, including stomach ulcer, Crohn's disease, diverticulitis, cancer of the stomach or bowel, or Ogilvie's syndrome.
- Tell your healthcare provider if you are pregnant or plan to become pregnant. Taking RELISTOR during pregnancy may cause opioid withdrawal symptoms in your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with RELISTOR.
- Taking RELISTOR while you are breastfeeding may cause opioid withdrawal in your baby. You should not breastfeed during treatment with RELISTOR. You and your healthcare provider should decide if you will take RELISTOR or breastfeed. You should not do both.
- Also, tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.
- In a clinical study, the most common side effects of RELISTOR tablets in people with long-lasting (chronic) pain that is not caused by cancer include: stomach-area (abdomen) pain, diarrhea, headache, swelling or a feeling of fullness or pressure in your abdomen, sweating, anxiety, muscle spasms, runny nose, and chills.

You are encouraged to report side effects of prescription drugs to FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For product information, adverse event reports, and product complaint reports, please contact: **Salix Product Information**

Call Center Phone: 1-800-321-4576 **Fax:** 1-510-595-8183

Email: salixmc@dlss.com

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Medication Guide.



Salix
PHARMACEUTICAL

400 Somerset Corporate Boulevard, Bridgewater, NJ 08807
Relistor is a trademark of Salix Pharmaceuticals or its affiliates.
All other trademarks are the property of their respective owners.
©2024 Salix Pharmaceuticals or its affiliates. REL.0103.USA.24

RELISTOR[®]
methylbuprenorphine hydrochloride
Tablets